

Supporting Information

Title: Repurposing of FDA-Approved Toremifene to Treat COVID-19 by blocking the spike glycoprotein and NSP14 of SARS-CoV-2

William R. Martin[⊥], and Feixiong Cheng^{⊥,∇,◇,*}

[⊥]Genomic Medicine Institute, Lerner Research Institute, Cleveland Clinic, Cleveland, OH 44195, USA

[∇]Department of Molecular Medicine, Cleveland Clinic Lerner College of Medicine, Case Western Reserve University, Cleveland, OH 44195, USA

[◇]Case Comprehensive Cancer Center, Case Western Reserve University School of Medicine, Cleveland, OH 44106, USA

*To whom correspondence should be addressed:

Feixiong Cheng, PhD

Lerner Research Institute, Cleveland Clinic

Tel: +1-216-444-7654; Fax: +1-216-636-0009

Email: chengf@ccf.org

Contents

Figure S1: a) NSP3 (PL-PRO) in the initial docked conformation b) NSP3 at 34.4 nanoseconds of simulation.

Figure S2: a) NSP4 in the initial docked conformation. b) NSP4 at 123.6 nanoseconds of simulation.

Figure S3: a) NSP5 (3CL-PRO) in its initial docked conformation. b) NSP5 at 284.2 nanoseconds of simulation.


Figure S4: a) NSP12 (RNA-Polymerase) in its initial docked conformation. b) NSP12 at 60.4 nanoseconds of simulation.

Figure S5: a) NSP15 in its initial docked conformation. b) NSP15 at 35.3 nanoseconds of simulation.

Figure S6: a) NSP16 in its initial docked conformation. b) NSP16 at 11.5 nanoseconds of simulation.

Figure S7: Root-Mean-Squared Deviation (RMSD) plot for all proteins simulated.

a



b





Figure S1: NSP3 (PL-PRO) in the initial docked conformation (a) and at 34.4 nanoseconds of simulation (b).

a



b





Figure S2: NSP4 in the initial docked conformation (a) and at 123.6 nanoseconds of simulation (b).

a



b




Figure S3: NSP5 (3CL-PRO) in its initial docked conformation (a) and at 284.2 nanoseconds of simulation (b).




Figure S4: NSP12 (RNA-Polymerase) in its initial docked conformation (a) and at 60.4 nanoseconds of simulation (b).




Figure S5: NSP15 in its initial docked conformation (a) and at 35.3 nanoseconds of simulation (b).




Figure S6: NSP16 in its initial docked conformation (a) and at 11.5 nanoseconds of simulation (b).




Figure S7: Root-Mean-Squared Deviation (RMSD) plot for all proteins simulated; each protein was submitted to 500 nanoseconds of molecular dynamics simulation.